Appendix Reducing Preventable Hospitalization and Disparity: Association With Local Health Department Mental Health Promotion Activities Chen et al.

Appendix Table 1. Prevention Quality Indicators (PQI) Using AHRQ Definition (Discharges, for Patients Aged ≥18 Years, With a Principal ICD-9-CM Diagnosis)

| PQIs | Inclusions | Exclusions ^a |
|-------------------------|---|--|
| PQ#1: Diabetes short- | 25010, 25011, 25012, 25013, | |
| term complications | 25020, 25021, 25022, 25023, | |
| diagnosis codes | 25030, 25031, 25032, 25033 | |
| PQ#3: Diabetes with | 25040, 25041, 25042, 25043, | |
| long-term | 25050, 25051, 25052, 25053, | |
| complications diagnosis | 25060, 25061, 25062, 25063, | |
| codes | 25070, 25071, 25072, 25073, | |
| | 25080, 25081, 25082, 25083, 25000, 25001, 25002, 25003 | |
| PQ#5: COPD or asthma | 25090, 25091, 25092, 25093 4910, 4911, 49120, 49121, | Cases with any-listed ICD-9-CM |
| in older adults | 4918, 4919, 4920, 4928, 484, | diagnosis codes for cystic fibrosis |
| in order addits | 4940, 4941, 496 | and anomalies of the respiratory |
| | 15.10, 15.11, 150 | system and obstetric admissions and |
| | | transfers from other institutions |
| | 49300, 49301, 49302, 49310, | |
| | 49311, 49312, 49320, 49321, | |
| | 49322, 49381, 49382, 49390, | |
| | 49391, 49392 | |
| | 4660, 490 | |
| PQ #7: Hypertension | 4010, 4019, 40200, 40210, | Cases with any-listed ICD-9-CM |
| diagnosis codes | 40290, 40300, 40310, 40390, | procedure codes for cardiac |
| | 40400, 40410, 40490 | procedure, with any listed ICD-9- |
| | | CM diagnosis codes of Stage I-IV kidney disease, only if accompanied |
| | | by any-listed ICD-9-CM procedure |
| | | codes for dialysis access |
| PQ #8: Heart failure | 39891, 4280, 4281, 42820, | Excludes cardiac procedure |
| diagnosis codes on or | 42821, 42822, 42823, 42830, | admissions, obstetric admissions, |
| after 10/01/2002 | 42831, 42832, 42833, 42840, | and transfers from other institutions |
| | 42841, 42842, 42843, 4289 | |
| PQ #13: Angina | 4111, 41181, 41189, 4130, | Cases with any-listed ICD-9-CM |
| diagnosis codes | 4131, 4139 | diagnosis codes for cardiac |
| DO 114 4 77 | 25002 25022 | procedure |
| PQ #14: Uncontrolled | 25002, 25003 | |
| diabetes without | | |
| mention of a short-term | | |
| or long-term | | |

Appendix Reducing Preventable Hospitalization and Disparity: Association With Local Health Department Mental Health Promotion Activities Chen et al.

| complication diagnosis | |
|------------------------|-----------------------------|
| codes | |
| PQ #15: Asthma | 49300, 49301, 49302, 49310, |
| diagnosis codes in | 49311, 49312, 49320, 49321, |
| younger adults | 49322, 49381, 49390, 49391, |
| | 49392 |

^aExclude cases: • transfer from a hospital (different facility) • transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF) • transfer from another healthcare facility. Source: Agency for Healthcare Research and Quality: Prevention Quality Indicators Technical Specifications.

https://www.qualityindicators.ahrq.gov/Modules/PQI_TechSpec_ICD10_v70.aspx.

AHRQ, the Agency for Healthcare Research and Quality; COPD, chronic obstructive pulmonary disease.

Appendix Reducing Preventable Hospitalization and Disparity: Association With Local Health Department Mental Health Promotion Activities Chen et al.

Appendix Table 2. ICD-9-CM Diagnosis Codes Used in Defining Study Conditions

| Description | ICD-9-CM codes |
|---|--|
| Mental and substance use disorders | |
| Adjustment disorders | 3090–3099 |
| Anxiety disorders | 30000–31383 |
| Attention-deficit, conduct, and | 31200–3149 |
| disruptive behavior disorders | |
| Elimination disorders | 3076–3077 |
| Other disorders usually diagnosed in | 3073, 30921, 31323, 31389, 3139, 29900, 29901, |
| infancy, childhood, or adolescence | 29910–29991, 30720–30723 |
| Eating disorders | 3071, 30750–30759 |
| Impulse control disorders, not | 31230–31239 |
| elsewhere classified | |
| Miscellaneous mental conditions | 30012–30019 |
| Psychogenic disorders | 3060–3069 |
| Sexual and gender identity disorders | 3020–3029, 30651 |
| Sleep disorders | 30740–30749 |
| Somatoform disorders | 30011, 3007, 30081, 30082, 30780–30789 |
| Bipolar disorders | 29600–29699 |
| Depressive disorders | 29620–29636, 3004, 311 |
| Personality disorders | 3010–3019 |
| Schizophrenia and other psychotic disorders | 29500–2989, 33392 |
| Alcohol-related disorders | 2910–2919, 30300–30393, 30500–30503, 3575, 4255, |
| | 5353, 5710–5713, 7903 |
| Drug abuse disorders | 2920–2929, 30400–30493, 30520–30593 |
| Drug dependence in pregnancy | 6483–64834, 6555–65553, 7795, 76072, 76073, 76075, |
| | 76071 |
| Mental disorders in pregnancy | 6484–64844 |
| Poisoning by opiates and related | 965000–96509 |
| narcotics | |
| Miscellaneous history codes related | V6542 |
| to substance abuse | |
| Miscellaneous history or injury | E95n.n, V110-V113, V118, V119, V154-V1549, |
| codes related to mental illness | V402, V403, V409, V663, V673, V7101, V7102, V7109, V790, V798, V799 |

Source: Coffey RM, Houchens R, Chu BC, Barrett M, Owens P, Stocks C, Vandivort-Warren R, Buck J, Emergency Department Use for Mental and Substance Use Disorders. Online August 23, 2010, U.S. Agency for Healthcare Research and Quality (AHRQ).